

Deep learning based depth map estimation of protoporphyrin IX in turbid media using dual wavelength excitation fluorescence: supplement

HINANO IMANISHI,¹ TAKAHIRO NISHIMURA,^{1,*} YU SHIMOJO,^{1,2,3}  AND KUNIO AWAZU^{1,4}

¹Graduate School of Engineering, Osaka University, Yamadaoka 2-1, Suita, Osaka 565-0871, Japan

²Graduate School of Medicine, Osaka Metropolitan University, Asahimachi 1-4-3, Abeno-ku, Osaka 545-8585, Japan

³Research Fellow of Japan Society for the Promotion of Science, Kojimachi 5-3-1, Chiyoda-ku, Tokyo 102-0083, Japan

⁴Global Center for Medical Engineering and Informatics, Osaka University, Yamadaoka 2-2, Suita, Osaka 565-0871, Japan

*nishimura-t@see.eng.osaka-u.ac.jp

This supplement published with Optica Publishing Group on 18 September 2023 by The Authors under the terms of the [Creative Commons Attribution 4.0 License](#) in the format provided by the authors and unedited. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Supplement DOI: <https://doi.org/10.6084/m9.figshare.24126957>

Parent Article DOI: <https://doi.org/10.1364/BOE.500022>

Deep learning based depth map estimation of protoporphyrin IX in turbid media using dual wavelength excitation fluorescence

Table S1. Optical Properties of Simulated Phantom. The optical properties were adjusted to ± 5 , ± 10 , $\pm 15\%$ relative to the baseline values.

		405 nm	505 nm	635 nm
Absorption coefficient [mm ⁻¹]	+15%	0.75	0.12	0.049
	+10%	0.72	0.11	0.047
	+5%	0.69	0.11	0.045
	Baseline	0.65	0.10	0.043
	-5%	0.62	0.099	0.041
	-10%	0.59	0.094	0.039
	-15%	0.56	0.088	0.037
Scattering coefficient [mm ⁻¹]	+15%	20	10.5	6.0
	+10%	19	10.0	5.7
	+5%	18	9.6	5.5
	Baseline	17	9.1	5.2
	-5%	16	8.7	5.0
	-10%	16	8.2	4.7
	-15%	15	7.8	4.4